

PATIENT INFORMATION / REFERRAL STATUS

Referral Status: ☐ New Referral ☐ Updated Order ☐ Order Renewal Date: _____

Patient Name: _____ DOB: _____

ICD-10 Code: _____ ICD-10 Description/Diagnosis: _____

Allergies: ☐ NKDA Allergies: _____ Weight: _____ ☐ lbs/☐ kg Height: _____

Patient Status: ☐ New to Therapy ☐ Continuing Therapy Last Treatment Date: _____ Next Due Date: _____

PROVIDER / PRACTICE INFORMATION

Ordering Provider: _____ Provider NPI: _____

Referring Practice Name: _____ Phone: _____ Fax: _____

Practice Address: _____ City: _____ State: _____ Zip: _____

Referral Coordinator Name: _____ Email: _____ Alternative Phone Number: _____

REFERRING PROVIDER COMMUNICATIONS

- ☐ I have reviewed the prescribing information and medication guide for Entyvio (vedolizumab).
- Avoid use of live vaccines during treatment. Prior to initiating therapy, complete all age-appropriate vaccinations according to current immunization guidelines.
 - Evaluate liver enzymes and bilirubin at baseline. Interrupt treatment if drug-induced liver injury is suspected. Instruct patients to seek immediate medical attention if they experience symptoms suggestive of hepatic dysfunction.
 - Evaluate patients for tuberculosis (TB) prior to initiating treatment. Do not administer to patients with active TB infection. Monitor patients for signs and symptoms of active TB during and after treatment.
 - Recommend evaluating patients for viral hepatitis prior to initiating treatment and treating them according to guidelines. Consider periodic evaluation of patients who are hepatitis B carriers for signs/symptoms of active hepatitis B infection.
 - Evaluate for active infection. Delay administration to patients with an active infection.

NURSING PROTOCOL COMMUNICATIONS

- ☒ Provide nursing care, vital signs, monitoring according to Memorial Outpatient Procedures. Establish/maintain IV access and administer medication as ordered. Remove peripheral IV access after infusion completion if applicable. Follow infusion-related/hypersensitivity reactions management according to MHS Outpatient Adverse Reaction Protocol available for review on at mhs.net/services/pharmacy/infusion-services/outpatient-infusion.
- ☒ Discharge/Follow-up instructions according to Memorial Outpatient Procedures.

LABORATORY ORDERS

- ☐ **Pregnancy, Urine for females of childbearing potential who have not undergone a hysterectomy:** ☐ Once ☐ Every Visit
- ☐ **CBC with Diff:** ☐ Once ☐ Every Visit ☐ Every ____ months
- ☐ **Comprehensive Metabolic Panel:** ☐ Once ☐ Every Visit ☐ Every ____ months
- ☐ **CRP:** ☐ Once ☐ Every Visit ☐ Every ____ months
- ☐ **Bilirubin fractionated:** ☐ Once ☐ Every Visit ☐ Every ____ months
- ☐ **Hep B surface antigen [HBsAg]:** ☐ Once ☐ Every Visit ☐ Every ____ months
- ☐ **Hep B surf. antibody quantitative:** ☐ Once ☐ Every Visit ☐ Every ____ months
- ☐ **Hep B core antibody, total [anti-HBc]:** ☐ Once ☐ Every Visit ☐ Every ____ months
- ☐ **Quantiferon (R) TB gold, draw site incubated:** ☐ Once ☐ Every ____ months



PRE-MEDICATION ORDERS (30-60 Minutes Prior to Therapy)

- ☐ Acetaminophen (Tylenol) 650 mg PO
- ☐ Diphenhydramine (Benadryl) ☐ 25 mg ☐ 50 mg ☐ PO ☐ IV **OR**
- ☐ Cetirizine (Zyrtec) or Loratadine (Claritin) 10 mg PO
- ☐ Methylprednisolone (Solu-Medrol) ☐ 40 mg ☐ 125 mg IV **OR**
- ☐ Dexamethasone (Decadron) ☐ 8 mg ☐ 20 mg PO
- ☐ Other: _____ Dose: _____ Route: _____ Frequency/Timing: _____

THERAPY PLAN**Medication Name:** Vedolizumab (Entyvio)**Induction Dose:**

- ☐ 300 mg in 250 mL 0.9% NS IV infusion over 30 minutes on Week 0, Week 2 and Week 6

Maintenance Dose:

- ☐ 300 mg in 250 mL 0.9% NS IV infusion over 30 minutes every 8 Weeks thereafter.
- ☐ 300 mg in 250 mL 0.9% NS IV infusion over 30 minutes on every _____ Weeks.

Route: ☒ IV****Diluent/Volume/Concentration/Special tubing/Filters will be in accordance with the product package insert.****

- ☒ Flush with 0.9% sodium chloride at completion per protocol or medication-specific instructions

Additional Administration Instructions:

Do not dilute or infuse through the same intravenous line with other solutions. Monitor for hypersensitivity reactions.

Note to Pharmacy/Comments:

Refills: ☐ Zero ☐ for 12 months ☐ Other: _____**(if not indicated, order will expire one year from date signed)**_____
*Provider Name (Print)*_____
*Provider Signature*_____
Date

Adverse Reaction Management Protocol (Adult)

Observe patient for infusion related and hypersensitivity reactions such as fever, chills, rigors, pruritus, rash, cough, sneezing, throat irritation, nausea, vomiting.

If reaction occurs:

- Stop infusion and assess patient.
- Maintain or establish vascular access if needed
- **Administer emergency medication(s) according to symptoms:**
 - ☒ Acetaminophen 650 mg PO once PRN headache, pain, fever >100.4F, chills or rigors.
 - ☒ Diphenhydramine 50 mg IV once PRN itching, allergies, infusion reaction, hives, pruritic and other nonspecific symptoms of allergic reaction **OR**
 - ☒ Diphenhydramine 50 mg IM once PRN itching, allergies, infusion reaction, hives, pruritic and other nonspecific symptoms of allergic reaction (if no IV access)
 - ☒ Dexamethasone 10 mg IV once PRN shortness of breath or wheezing **OR**
 - ☒ Dexamethasone 10 mg IM once PRN shortness of breath or wheezing (if no IV access)
 - ☒ Ondansetron 4 mg IV once PRN nausea, vomiting **OR**
 - ☒ Ondansetron 4 mg IM once PRN nausea, vomiting (if no IV access)
- May re-start therapy if appropriate when symptoms resolve. Resume infusion at 50% of the previous rate and increase per manufacturer's guidelines.

If a severe allergic/anaphylactic reaction occurs

- Symptoms are rapidly progressing or continuing after administration of PRN medications and/or signs and symptoms of severe allergic/anaphylactic reaction (angioedema, swelling of the mouth, tongue, lips, or airway, dyspnea, bronchospasm with or without hypotension or hypertension)
 - ☒ Notify the Rapid Response / Rescue Alert Team / Blue Alert / 911.
 - ☒ Initiate BLS/ Cardiopulmonary resuscitation if necessary.
 - ☒ Administer Epinephrine 0.3 mg intramuscularly, every 5 MIN PRN rapidly progressing or continuing after administration of PRN medication or signs and symptoms of severe allergic/anaphylactic reaction. Administer every 5-15 minutes as needed preferably in the outer thigh.
 - ☒ Place the patient in a recumbent position, elevate lower extremities.
 - ☒ Continuously monitor vital signs (blood pressure, pulse oximetry, and heart rate).
 - ☒ Contact and notify the referring provider on the day of occurrence once patient is stabilized.
 - ☒ Document reaction in the medical record and complete an incident report once patient is stabilized.